

RETARDENT NC THINNER Product No. **TN122**

RETARDANT NC THINNER is a mixture of glycol solvents. It is recommended to use in a certain proportion with NC thinner (5%) to prevent milky effect or ruff finish for all Nitrocellulose varnish products especially during summer and higher temperature than normal, due to its slow evaporation.

SPECIFICATIONS

Properties

Color	: Clear
Specific gravity	: 0.92 Kg/lit [± 0.025]
Initial Boiling Point	: 150°C
Non Aromatics	: 100%
Flash point	: 21°C

APPLICATION METHODS

All RITVER thinners listed are to be used in conjunction with the corresponding products and systems intended for, or any Technical guidance required may be obtained by the end user.

Advantage: Improves film smoothness of lacquers when applied either by spray or dipping thus extending excellent finish.

Lacquer driers without blushing, resin blush, pin holing and other surface defects.

No orange peel developed after the evaporation of NC Thinner, and therefore increase the tensile strength of the film along with its gloss.

NC Retardent Thinner has a certain action preventing milky effect of the surface.

PACKING & STORAGE

Available in 1Ltr, 5Ltr, 5USG, 10x5USG.

Store in a cool and well ventilated place. Keep away from direct sunlight. 18 months in unopened container, stored in a cool and dry condition at 25°C.

RETARDENT NC THINNER Product No. **TN122**

HANDLING

Disposal: As per the guidance and legislations of the local Authority e.g., by controlled landfill. In case of doubt, consult local authority. Do not empty into drains, sewers or other water courses.

Flash Point: Individual flash points are mentioned in the table along with other details and a prior knowledge of the flash points is a pre requisite for the safe handling of the **RITVER thinners**.

SAFETY PRECAUTION

First Aid:

Eyes In the event of accidental splashes, flush eyes with warm water immediately and obtain medical advice.

Skin Wash skin thoroughly with soap and water or approved Industrial cleaner. DO NOT USE solvent or thinners.

Inhalation Remove to fresh air, loosen collar and keep patient rested.

Ingestion In case of accidental ingestion DO NOT INDUCES VOMITING. Obtain immediate medical attention.